

Traszka karpacka *Lissotriton montandoni* (Boulenger, 1880) w pokarmie pstrąga potokowego *Salmo trutta* m. *fario* L.

Montandon's newt *Lissotriton montandoni* (Boulenger, 1880)
in the brown trout *Salmo trutta* m. *fario* L. diet.

ANETA BYLAK¹, KRZYSZTOF KUKUŁA¹

¹Katedra Biologii Środowiska, Uniwersytet Rzeszowski
35-310 Rzeszów, ul. Prof. S. Pigoń 6
e-mail: abyлак@univ.rzeszow.pl, kkukula@univ.rzeszow.pl

Słowa kluczowe: pstrąg potokowy, pokarm, stawy bobrowe, Bieszczady Wysokie.

Summary

The diet of large (*TL* 37.3 cm) brown trout *Salmo trutta* m. *fario* L. from a beaver pond in the Bieszczady National Park was studied. The trout diet was mainly composed by Montandon's newts *Lissotriton montandoni*. Fish and other vertebrates were not observed in trout's stomach. The beavers have dammed the mountain stream and created the new habitat with stagnant water. Deep, well-oxygenated beaver ponds are suitable for large fish and also for newts, during its mating season. Thus, adult newts were a prey item in the brown trout diet.